

## ENTATION PAGE

Form Approved  
OMB No. 0704-0188

AD-A232 837

estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering the collection of information. Send comments regarding this burden estimate or any other aspect of this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Ave, Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

## PORT DATE

January 1991

## 3. REPORT TYPE AND DATES COVERED

Final

11/1/87 - 10/31/90

## 4. TITLE AND SUBTITLE

Atrane and Quadroped Frameworks in Early Transition metal  
and Silicon Gro Stereochemistry: Monomers and Oligomers

## 5. FUNDING NUMBERS

2303/B2 61102F

## 6. AUTHOR(S)

J. G. Verkade

## 7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)

Iowa State University  
Department of Chemistry  
Ames, IA 50011

8. PERFORMING ORGANIZATION  
REPORT NUMBER

AFOSR-TR- 91 0174

## 9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)

AFOSR/NC Bd 410  
Bolling AFB, DC 20332-6448

10. SPONSORING/MONITORING  
AGENCY REPORT NUMBER

AFOSR-88-0081

## 11. SUPPLEMENTARY NOTES

DTIC  
ELECTE  
MAR 13 1991  
S B D

## 12a. DISTRIBUTION/AVAILABILITY STATEMENT

Approved for public release; distribution is unlimited

## 12b. DISTRIBUTION CODE

## 13. ABSTRACT (Maximum 200 words)

An exploration of the chemistry of silatranes and azasilatranes has been carried out. The first isolation and structural characterization of silatrane adducts in which the exocyclic axial oxygen behaves as a Lewis base site was accomplished. Azasilatrane phosphine-nitrogen complexes were prepared with Lewis acid sites on the complex bonding to the axial oxygen. An axial silatranium cation was prepared. The range of axial-substituents for azasilatranes was explored and found to be limited to H, OR, OAr, R, and Ar. Attempts to obtain pentacoordination by nitrogen by replacing the axial hydrogen by azide using trimethylsilylazide resulted instead in trimethylsilyl substitution of the equatorial nitrogens and an azide salt. Pyrolysis of the azide salt afforded the pentacoordinated azide. Nitrogen pentacoordination was also obtained from a thioisocyanide salt. A number of other equatorial nitrogen substituted azasilatranes were prepared. Attempts to couple azasilatrane to a Si-Si bonded dimer were unsuccessful. Azasilatrane-based polymers were prepared and characterized. The azasilatrane studies were extended to azastannatranes.

## 14. SUBJECT TERMS

## 15. NUMBER OF PAGES

15

## 16. PRICE CODE

17. SECURITY CLASSIFICATION  
OF REPORT

UNCLASSIFIED

18. SECURITY CLASSIFICATION  
OF THIS PAGE

UNCLASSIFIED

19. SECURITY CLASSIFICATION  
OF ABSTRACT

UNCLASSIFIED

## 20. LIMITATION OF ABSTRACT

SAR

**Final Technical Report**

**J. G. Verkade**

**AFOSR-88-0081**

**November 1, 1987 - October 30, 1990**


Approved for public release;  
distribution unlimited.

**91 3 11 136**


It should be noted at the outset that the PI's research described in this section was supported by one three-year AFOSR grant which expired October 31, 1990. A preproposal for renewal was approved by the AFOSR June 12, 1990 and the full proposal based on the preproposal was mailed July 20, 1990. On July 30 the PI was informed by telephone by the AFOSR that although this new departure in our research program had been unusually productive, recent fiscal cutbacks and changes in programmatic priorities necessitated that the full proposal be declined without review.

The primary objectives of the AFOSR grant were:

A. Exploration of the chemistry of silatranes such as 1 and azasilatranes such as 2.



1 Z = R, OR; E = O  
2 Z = R, OR; E = NR



3

B. Synthesis of silatranes in which silicon is pentacoordinated by nitrogens as in 3.

C. Synthesis of oligomers/polymers containing azasilatrane repeating units.

All three of these objectives were achieved.

Though beyond the scope of the AFOSR grant, the eventual goal towards which our research was directed was the synthesis of polymers such as 4 and 5 which might serve as precursors to essentially pure silicon nitride, with minimum contamination by silicon carbide, owing to the robustness of the chelated hypercoordinated silicon and the presence of only Si-N bonds in the azasilatrane cage.

What follows is a summary of the work associated with the objectives A-C given above.




Distribution/  
Availability Codes


Dist Avail and/or  
Special

Dist

A-1



4




5

### A. Exploration of the chemistry of silatranes and azasilatranes

In order to construct polymers of types 4 and 5, it was necessary to gain some knowledge of the reactivities (nucleophilicities) of the exocyclic axial, the equatorial, and the endocyclic axial substituents on the silicon in 1 and 2. The chemistries of these reactivity sites are now addressed in turn.

#### 1. Reactivity of the axial substituent in silatranes and azasilatranes

The equatorial and axial oxygens in alkoxy silatranes such as 6 and 7 might be




expected to display increased basicity owing to electron induction into silicon by the *trans* axial tertiary amino nitrogen. We recently reported the first isolation and structural characterization of silatrane adducts in which the exocyclic axial oxygen behaves as the Lewis basic site.<sup>1</sup> Thus silatranes 6 and 7 are fully covalently bonded and hydrogen bonded to their Lewis acid components in the structures of 8(BF<sub>4</sub>) and 9, respectively. The SiN<sub>ax</sub> bond length decreases considerably (~0.5 Å) from 7 to 8(BF<sub>4</sub>) (1.965(5) Å) and the

latter is the shortest such bond distance reported for any silatrane. The O(H)O distance in **9** (2.489 Å) is the shortest reported for an unsymmetrical hydrogen bond. The SiO<sub>ax</sub> distance increases by a total of 0.17 Å in the order **7** < **9** < **8**(BF<sub>4</sub>).


Interestingly, cation **8** is the first reported example of a silatrane having an ideal TBP symmetry. Thus all others that have been structured contain silicon atoms that are at least ~0.1 Å above the O<sub>eq</sub> plane. In this connection we have shown<sup>1</sup> that a prediction<sup>2</sup> that an ideally TBP silatrane should possess an Si-N<sub>ax</sub> distance of 1.83 Å is incorrect by more than 0.1 Å.

Measurement of phenol ν(OH) shifts reveals the basicity order (Me<sub>3</sub>Si)<sub>2</sub>O < Si(OR)<sub>4</sub> < alkoxy silatranes ≤ Me<sub>3</sub>SiOMe < Et<sub>2</sub>O. While this trend places the electron inductive ability of the silatranyl group ahead of (RO)<sub>3</sub>Si, it is below that of an Et group.


Until we began the work supported by the AFOSR grant, there were only four azasilatranes (**2**) reported and very little was known about them. Structural evidence is accumulating, however, that these compounds possess Si-N<sub>ax</sub> bonds at least as robust as those in silatranes (**1**).<sup>3,4</sup> This is somewhat unexpected in view of the greater electronegativity of the equatorial oxygen substituents in **1** than in **2**. Thus the Si-N<sub>ax</sub> bond lengths in **10**,<sup>3</sup> **11**<sup>4</sup> and **12**<sup>4</sup> are very comparable to that in **7**.<sup>1</sup> That the ethoxy oxygen




in these azasilatranes also displays Lewis base character is evident in complexes **13** and **14** which can be converted to **15** and **16**, respectively in the presence of CO.



We carried out several interesting substitution reactions at the axial position. In reaction 2 the synthesis of the novel silatranium cation 18 is realized under very mild




conditions.<sup>5</sup> Compound 18 is, however, very resistant to nucleophilic attack by  $\text{F}^-$ ,  $\text{H}^-$  and  $\text{OR}^-$ .




We converted azasilatran 19 to 20<sup>3</sup> and 21,<sup>6</sup> which we planned to use as precursors to amido azasilatranes. Unlike silanes,<sup>7</sup> 19 is unreactive to Si-H proton substitution by


amide.<sup>3</sup> Unfortunately the range of Z groups in **22** formed via reaction 5 seems to be restricted to Z = H, OR, OAr, R and Ar.<sup>3</sup>



Attempts to synthesize the novel structure **23**, which because of its zero dipole dipole moment, may be a volatile MOCVD precursor for SiN, were unsuccessful.<sup>8</sup>




The above studies were being extended to azastannatranes such as **24-27**, a class of compounds we synthesized for the first time.<sup>6</sup> A most curious observation in the crystal structure determination of azastannatrane **26** is that the two independent molecules in the unit cell possess substantially different Sn-N<sub>ax</sub> bond lengths of 2.380(2) and 2.453(2) Å.<sup>9</sup> Two different <sup>119</sup>Sn chemical shifts in the solid state have also been observed.<sup>9</sup> The most reasonable rationale for this observation is differences in crystal packing forces. This rationale has also been advanced<sup>10</sup> to account for the observation that silatranes in the solid state, such as those we have structured, frequently exhibit shorter Si-N<sub>ax</sub> distances than are found calculationaly or in the gas phase by electron diffraction.



## 2. Reactivity of the equatorial substituent in silatranes and azasilatranes

Discussed in this section are addition of  $\text{H}^+$  and  $\text{R}^+$  to an  $\text{N}_{\text{eq}}$  site, displacement of the  $\text{N}_{\text{eq}}\text{-H}$  hydrogen by R, and solvolytic cleavage of  $\text{Si-N}_{\text{eq}}$  bonds.


In a recent communication<sup>11</sup> we reported the synthesis of 28, the first stable representative of incipient proton-assisted equatorial bond cleavage of a five-coordinate intermediate to be isolated.<sup>11</sup> Reaction 8 actually constituted an attempt to substitute the




axial proton of 19 by azide. Instead,  $\text{Me}_3\text{Si}^+$  attacks  $\text{N}_{\text{eq}}$  sites, liberating the protonating agent  $\text{HN}_3$ . Interestingly compound 28 co-crystallized with 19 in the unit cell of 29. The molecular structure of 29<sup>11</sup> as well as its analogue 33,<sup>12</sup> determined by X-ray means, revealed that in each case the  $\text{Si-N}_{\text{eq}}\text{H}_2$  length is ca. 10% longer than the two  $\text{Si-N}_{\text{eq}}\text{H}$  bonds in this cation. By analogous reactions of azasilatranes with  $\text{MeOTf}$  and  $\text{Me}_3\text{SiOTf}$ , the new salts 34-39 were synthesized.<sup>13</sup>

We carried out a variety of substitution reactions at the  $\text{N}_{\text{eq}}\text{-H}$  function in azasilatranes as exemplified by reaction 9 and Scheme I.<sup>3,14</sup> Though the sterically elongated  $\text{Si-N}_{\text{ax}}$  distance in 51 (2.775(7) Å) is the longest ever recorded in an azasilatrane,





it is 24% shorter than the sum of the van der Waals radii, suggesting some degree of silicon interaction with the nearly planar N<sub>ax</sub>.<sup>14</sup>




Solvolysis experiments on azasilatranes with MeOH revealed that the Si-N<sub>eq</sub> bond is more sensitive than the axial Si-H, Si-OEt or Si-CH<sub>3</sub> bond.<sup>15</sup> In Schemes II-V it is seen that a) solvolysis of the substituent on N<sub>eq</sub> may or may not precede Si-N<sub>eq</sub> bond cleavage and b) monocyclic solvolysis intermediates are sometimes detected.

The rate and pathway of methanolysis appear to be strongly dependent on the strength of the N<sub>eq</sub>-substituent bond and on the steric crowding these groups experience with the exocyclic axial group.<sup>15</sup>


### 3. Reactivity of the endocyclic axial substituent in azasilatranes

Although substantial Si-N<sub>ax</sub> bond strength in azasilatranes is afforded by the presence of the three chelating five-membered rings, weakening of this bond has been seen to occur in the previous section by steric crowding (e.g., 51) and cleavage can take


Scheme I




Scheme II




## Scheme III




## Scheme IV






Scheme V




place by solvolytic cleavage of one of the Si-N<sub>eq</sub> bonds (e.g., 54). The weakened Si-N<sub>ax</sub> interaction in 51 allowed us to synthesize 60<sup>14</sup> in which the Si-N<sub>ax</sub> distance (3.561 Å)<sup>15</sup> has increased from 2.775 Å in 51 to being close to the van der Waals radius (3.65 Å). The BH<sub>3</sub> adduct of 51 (61) is currently being structured.<sup>16</sup>

The reaction in which equatorially silylated 38 is formed also affords the axially silylated isomer 62 in an equilibrium which depends upon solvent polarity and temperature.<sup>12</sup> Arguments substantiating the possibility of a hypercoordinate nitrogen in 62 have been brought forward.<sup>13</sup>




### B. Synthesis of silatranes pentacoordinated by nitrogen ligands


Although rational efforts to synthesize compounds of this type have thus far not been successful (See Section II. A. 1.; note however, 27) compound 29 upon heating under vacuum yields 63.<sup>12</sup> A similar reaction (also involving disproportionation, however)



produces 64 from 33.<sup>12</sup> These are the first reported examples of silatrane-type systems




containing five nitrogens coordinated to the silicon. We also have preliminary evidence that **65** is formed in reaction 12.<sup>8</sup>



### C. Synthesis of oligomers/polymers containing azasilatrane units


Two azasilatrane-based oligomers/polymers were partially characterized. The second step in Scheme VI gives rise to a small amount of a sublimate which may be **66**. Further heating of the residue gives a solid **67** which has appreciable solubility only in DMSO.

Scheme VI






The alternate routes in Scheme VII give a solid **68** which, encouragingly, is chloroform soluble. The <sup>29</sup>Si NMR spectrum of **68** displays two broad peaks indicative of four-coordinate (-25 ppm) and five-coordinate (-77 ppm) silicon. The <sup>1</sup>H NMR spectrum exhibits N(CH<sub>3</sub>)<sub>2</sub> peaks but no detectable NH resonance, suggesting that all three N<sub>eq</sub>-H

## Scheme VII



protons in **19** react with  $(\text{Me}_2\text{N})_3\text{SiH}$  possibly giving rise to "star" polymer in which the equatorial nitrogens in the triangular  $\text{SiN}_3$  moieties of the azasilatrane units are bridged by  $\text{HSiNMe}_2$  groups.

Polymers **69-71** shown in reactions 13-15, respectively, formed under similar



conditions and were studied by NMR spectroscopy in both solution and the solid state.  $^{29}\text{Si}$  NMR studies confirm the presence of both four and five-coordinate Si in all cases. The absence of  $\text{NMe}_2$  and  $\text{NH}$  functions in **69** was inferred from DEPT- $^{13}\text{C}$  and DEPT- $^1\text{H}$  spectra, respectively. While  $\text{NH}$  groups are not apparent in **70**,  $\text{NMe}_2$  groups do appear in both the  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectra. Using  $^1\text{H}$  -  $^{29}\text{Si}$  polarization transfer techniques, all the  $\text{H-Si}$  groups in **71** appear in the four-coordinate  $^{29}\text{Si}$  region, while the  $\text{MeSi}$  moieties (originating from the starting atrane **47**) resonate in the five-coordinate  $^{29}\text{Si}$  region. In

contrast to **69** and **70**, **71** possesses NH groups as judged from the  $^1\text{H}$  NMR and IR spectra. The spectroscopic data are consistent with a linear polymer wherein one of the three NH functions on the silatrane **47** does not react.

**D. References** (Asterisked references represent publications supported by the AFOSR grant.)

- \*(1) Garant, R. J.; Daniels, L. M.; Das, S. K.; Janakiraman, M. N.; Jacobson, R. A.; Verkade, J. G., provisionally accepted by *J. Am. Chem. Soc.*
- (2) Sidorkin, S. F.; Pestunovich, V. A.; Voronkov, M. G. *Russ. Chem. Rev.* (Engl. Transl.) 1980, 49, 414.
- \*(3) Gudat, D.; Verkade, J. G. *Organometallics* 1989, 8, 2772.
- \*(4) Gudat, D.; Daniels, L. M.; Verkade, J. G. *Organometallics* 1990, 10, 1664.
- \*(5) Garant, R. J.; Plass, W.; Verkade, J. G., manuscript in progress.
- \*(6) Plass, W.; Verkade, J. G., to be published.
- (7) Pawlenko, S. In *Houben-Weyl, Bd XIII/5*; Georg Thieme Verlag: Stuttgart, 1980; S. 227f.
- \*(8) Wan, Y.-J.; Verkade, J. G., to be published.
- \*(9) Plass, W.; Daniels, L. M.; Verkade, to be published.
- (10) Davis, I., 1990, private communication.
- \*(11) Woning, J.; Daniels, L. M.; Verkade, J. G. *J. Am. Chem. Soc.* 1990, 112, 4601.
- \*(12) Woning, J.; Verkade, J. G., *Organometallics*, provisionally accepted.
- \*(13) Verkade, J. G.; Woning, J., submitted to *J. Am. Chem. Soc.*
- \*(14) Gudat, D.; Daniels, L. M.; Verkade, J. G. *J. Am. Chem. Soc.* 1989, 111, 8520.
- \*(15) Gudat, D.; Verkade, J. G. *Organometallics* 1990, 9, 2172.
- \*(16) Gudat, D.; Woning, J.; Daniels, L. M.; Verkade, J. G., to be published.
- \*(17) Gudat, D.; Lensink, C.; Schmidt, H.; Xi, S.-K.; Verkade, J. G. *Phosphorus, Sulfur, Silicon* 1988, 41, 21. Invited lecture, IRIS Meeting, Amherst, MA, June, 1988.